



Skin needling as a treatment for acne scarring: An up-to-date review of the literature[☆]

Adam G. Harris, MBChB, Catherine Naidoo, MBBS, Dedee F. Murrell, MA, BMBCh, MD, FACD, FRCP^{*}

Department of Dermatology, St George Hospital, Sydney, Australia
University of New South Wales, Sydney, Australia

ARTICLE INFO

Article history:

Received 28 January 2015

Received in revised form 9 March 2015

Accepted 18 March 2015

Keywords:

Acne scarring
Dermaroller
Microneedling
Micro-needling
Percutaneous collagen induction
Skin needling

ABSTRACT

Background: Skin needling is a technique used to improve the appearance of acne scarring.

Objective: To comprehensively review the medical literature regarding skin needling as a treatment for acne scarring.

Methods: A literature search was performed using the PubMed, Medline, and Embase databases, in addition to reviewing the bibliographies of relevant articles.

Results: Ten studies presented patients treated with skin needling alone, while eight studies discussed skin needling in combination with other treatments for acne scarring. All studies showed improvements in scarring after needling, with 12 reporting statistical significance. The median number of treatments when needling was used alone was three, the median duration between treatments was 4 weeks, and the median needle length used was 1.5 mm. Reported adverse events were infrequent and included post-inflammatory hyperpigmentation, "tram track" scarring, acne, and milia. There were no reports of bacterial infections.

Limitations: The studies reviewed were heterogeneous in design and of variable validity, with some not reporting statistical significance.

Conclusion: There is moderate evidence to suggest that skin needling is beneficial and safe for the treatment of acne scarring. However, double-blinded, randomized controlled trials are required to make more definitive conclusions.

© 2015 The Authors. Published by Elsevier Inc. on behalf of Women's Dermatologic Society. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Skin needling is a technique predominantly used to improve the appearance of cutaneous scarring and photodamage (Fig. 1). Fine needles puncture the skin, resulting in increased dermal elastin and collagen, collagen remodeling, and thickening of the epidermis and dermis (Aust et al., 2008a, 2010a, 2010b, 2011; Fabbrocini et al., 2011a; Fernandes, 2005; Fernandes and Signorini, 2008; Kim et al., 2011; Park et al., 2012; Schwarz and Laaff, 2011). Additionally, skin needling creates small channels, which increase the absorption of topically applied preparations (Badran et al., 2009; Kalluri et al., 2011), a property which has been used in various dermatological treatments (Bencini et al., 2012; Budamakuntla et al., 2013; Clementoni et al., 2010; Fabbrocini et al., 2011b, 2014a; Kang et al., 2008; Torezan et al., 2013; Yoon et al., 2008, 2010).

Skin needling for cutaneous scarring was introduced into the medical literature in 1997, when tattooing without pigment was used to abrade facial scars, improving their quality and color (Camirand and Doucet, 1997). In 1998, Desmond Fernandes, a plastic surgeon from South Africa, designed a hand-held device composed of a rolling barrel

with multiple protruding needles and used it for a technique he termed "percutaneous collagen induction." He later published his experience using this technique for various dermatological conditions, including acne scarring (Fernandes, 2005; Fernandes and Signorini, 2008).

Both manual and electronic hand-held skin needling devices are now widely available as low-cost therapies for the treatment of acne scarring (Doddaballapur, 2009; Dogra et al., 2014), yet, to our knowledge, there has been no dedicated review of the medical literature regarding this topic.

Objective

The objective of this study was to comprehensively review the medical literature regarding skin needling as a treatment for acne scarring.

Material and methods

A literature search was performed using the PubMed, Medline 1946-to-present, and Embase classic plus Embase 1947-to-present databases. Search terms included 'skin needling', 'microneedling', 'needle dermabrasion', 'tattooing without pigment', 'dry tattooing', 'percutaneous collagen induction therapy', and 'dermaroller'. All terms are used in the medical literature to describe skin needling and were combined with the term acne. The search was performed without limitations or

[☆] Conflicts of interest: The authors state no conflicts of interest.

^{*} Corresponding author.

E-mail address: d.murrell@unsw.edu.au (D.F. Murrell).



Fig. 1. An improvement in acne scarring and photodamage in a patient treated with one session of skin needling.

language restrictions. All bibliographies within the relevant articles were reviewed for their relevance.

Results

Nineteen articles included patients with acne scarring treated with skin needling. Included were nine prospective observational studies (Beretta et al., 2008; Dogra et al., 2014; Fabbrocini et al., 2009, 2014b; Garg and Baveja, 2014; Kang et al., 2009; Kim, 2008; Majid, 2009; Schwarz and Laaff, 2011), nine prospective controlled studies (Alam et al., 2014; Fabbrocini et al., 2011c; Gadkari and Nayak, 2014; Nofal et al., 2014; Leheta et al., 2011, 2014a, 2014b; Mohammed, 2013; Sharad, 2011), and one case report (Pahwa et al., 2012). Additionally, one prospective observational study likely contained patients with acne scarring (Aust et al., 2008a), but the method was unclear. Eighteen articles were in English and one was in Italian, which was translated. All studies were critically evaluated and are summarized in tables that can be found in the Supplementary Material.

Efficacy

The majority of the studies reviewed included an objective scar scoring system to measure efficacy, most commonly the grading systems designed by Goodman and Baron (Goodman and Baron, 2006a, 2006b). Others used subjective scar scoring scales, subjective improvement scales, and patient satisfaction scales. For the purpose of this review, if an objective scar scoring scale was not used, the results of a subjective scale were used to evaluate efficacy.

Studies evaluating the efficacy of skin needling alone as a treatment for acne scarring

Six studies measured the efficacy of skin needling alone as a treatment for acne scarring (Alam et al., 2014; Beretta et al., 2008; Camirand and Doucet, 1997; Fabbrocini et al., 2009, 2014b; Majid, 2009). All studies showed improvements in scar severity scores compared to baseline, with statistical significance reported in all except one. One study was a randomized placebo controlled trial (Alam et al., 2014); the remaining five were prospective observational trials, which are subject to a high risk of selection bias and inherently lower validity.

The first study published included 20 patients assessed before and after treatment by two physicians, blinded to pretreatment scores using the Goodman and Baron grading systems (Beretta et al., 2008).

After treatment, there was a statistically significant improvement in scores, but it was unclear how this was calculated.

A similar study included 37 patients with facial scarring, 32 of whom had acne scarring (Majid, 2009). A single dermatologist consecutively graded patients using the qualitative Goodman and Baron grading system. Twenty-seven of the 31 patients with acne scarring who completed the study had improved scar grades, but there was no statistical analysis of the results.

An Italian group published two further observational studies (Fabbrocini et al., 2009, 2014b). The first (2009) included 32 patients consecutively graded by the same dermatologist using the qualitative Goodman and Baron grading system. After treatment, there was a statistically significant reduction in the mean severity grading. A second, larger study (2014b) included 60 patients whose scar severity was evaluated from photographs using the Global Aesthetic Improvement Scale. After treatment, there was also a statistically significant reduction in severity grading.

A recent uncontrolled study from India (Dogra et al., 2014) included 36 patients assessed with the acne scar assessment tool described by Peterson et al. (2011). After treatment, there was a statistically significant improvement in the mean scar grading from 11.73 to 6.5, although six patients did not complete the study, five because of treatment-related complications including severe post-inflammatory hyperpigmentation and tram-track scarring.

The most rigorous study was a placebo-controlled, split-face trial from the United States, which included 20 participants randomized to receive either skin needling with topical anesthetic or topical anesthetic



Fig. 2. Development of erythema, crusting, and pustules in a patient one day after skin needling.

alone to either side of the face (Alam et al., 2014). Two dermatologists blinded to the intervention graded standardized photographs with the quantitative Goodman and Baron grading system. At 6 months post-treatment, there was a statistically significant decrease in the mean grade of the skin-needled side compared to the placebo side, with a mean difference in scores of 3.4 compared to 0.4. Five patients dropped out before the study protocol was initiated. There was no blinding of the patients.

An additional study included a prospective sub-analysis of 15 patients with “scars and stretch marks,” but it was not clear if the group contained patients with acne scarring (Aust et al., 2008a). Patients were a part of a larger group included in a retrospective analysis of skin needling used for a variety of conditions including acne scarring. The subgroup demonstrated a statistically significant improvement based on the Vancouver Scar Scale (Baryza, 1995) and Observer Scar Assessment Scale as assessed by two independent observers.

Studies evaluating the efficacy of skin needling compared to other methods of treatment for acne scarring

A total of three studies compared skin needling to other methods of treatment for acne scarring.

An Egyptian group (Leheta et al., 2011) randomized 30 participants to receive either skin needling or the focal application of 100% trichloroacetic acid (TCA) using the CROSS (chemical reconstruction of skin scars) method (Lee et al., 2002). A blinded dermatologist scored a 68% mean improvement in the needling group and a 75% improvement in the TCA group, with no statistically significant difference between groups. Participants were not blinded and three participants in the TCA group dropped out and were not included in the analysis on an intention-to-treat basis.

An Italian group compared skin needling to skin needling combined with the topical application of platelet-rich plasma (PRP) in a split-face trial of 12 patients (Fabbrocini et al., 2011c). After treatment, all scores were reduced but the PRP group had an overall mean lower severity score. The statistical significance of this result was not reported. There was also no mention of blinding, either of the investigator or the patient, or randomization to which side received each treatment.

A study from India sequentially enrolled 30 patients and compared skin needling to the combination of skin needling alternating with 35% glycolic acid (GA) peels (Sharad, 2011). Scars were graded by the treating dermatologist using the Echelle d'Evaluation clinique des Cicatrices d'acne classification (Dreno et al., 2007). The combination treatment resulted in a statistically significant greater mean improvement of 63% compared to 31%. The patients and the assessors were not blinded.

Studies evaluating the efficacy of skin needling in conjunction with other treatments for acne scarring

Skin needling was used in eight studies to increase the penetration of topically applied preparations and alongside other treatments to synergistically improve efficacy.

An Egyptian group randomly assigned 24 patients to receive skin needling and 20% TCA or deep skin peeling using 60% phenol (Leheta et al., 2014a). Live assessment by a dermatologist blinded to the intervention gave statistically significant mean improvements in scores of 70% in the combination group and 75% in the deep phenol peel group with no statistically significant difference between two. A second study by the group randomly assigned 39 patients to skin needling and 20% TCA (group I), fractional thermolysis (group II), or a combination of both treatments (group III) (Leheta et al., 2014b). The same blinded dermatologist scored patients and gave an improvement in mean severity scores of 60%, 62%, and 78%, respectively. There was no statistically significant difference between groups I and II, but there was between groups I and II versus group III. One patient from group I was lost to follow-up and was included in the analysis on an

intention-to-treat basis. Two patients from each group in the first study dropped out and were not included in the analysis. In both studies the patients were not blinded.

Another study combined skin needling, subcision, and 15% TCA in 50 patients (Garg and Baveja, 2014). Photographs were graded by the same nontreating physician using the qualitative Goodman and Baron grading system. Scar grades improved in all patients and although the process of statistical analysis was mentioned, it did not appear to be performed and presented.

A Korean study (Kang et al., 2009) assessed the combination of the focal application of 100% TCA, skin needling using a 29-gauge needle, subcision (Orentreich and Orentreich, 1995), and fractional thermolysis in 35 patients. An independent physician scored patients using the acne severity scale described by Lipper and Perez (2006). Ten patients completed the study, with scores improving in all patients. Statistical significance of the results was not reported and there was a large loss to follow-up.

An Indian study compared skin needling combined with subcision to subcision combined with cryorolling (Gadkari and Nayak, 2014). Cryorolling consisted of dipping the needling device into liquid nitrogen immediately before the procedure. Thirty-seven patients were randomized to have both procedures to either side of their face and a blinded observer scored standardized photographs using the qualitative and quantitative Goodman and Baron grading systems. Both treatments resulted in statistically significant improvements in mean quantitative gradings of 57% in the cryoroller group and 40% in the needling group. The difference between the two was statistically significant. Seven patients dropped out of the study and it was unclear which group they were from and whether they were included with intention to treat.

Another Egyptian study compared 45 patients equally randomized into three groups to receive either skin needling combined with the topical application of PRP, the focal application of 100% TCA, or intralesional dermal injections of PRP (Nofal et al., 2014). Photographs were assessed by two blinded dermatologists using the qualitative Goodman and Baron grading system. All three treatments resulted in statistically significant improvements in scar grades with no difference between them.

Finally, two separate studies looked at the combination CO₂ laser and skin needling using a 26-gauge needle. The first included 35 patients who all had improvements in a 4-point improvement scale. There was no statistical analysis of the results (Kim, 2008). The second randomized 60 patients to compare laser and skin needling to laser alone (Mohammed, 2013). Patients were assessed by three independent observers blinded to the treatment using the quantitative Goodman and Baron grading system. After treatment there was a statistically significant improvement in scores in both groups with no statistically significant difference between them.

Histological changes

One study published histological sections before and after skin needling treatment of 10 patients with posttraumatic and acne scarring (Schwarz and Laaff, 2011). A blinded dermatologist and pathologist concluded that, in seven patients, there was a noticeable increase in elastin correlating with the depth of needle penetration. Increases in collagen and dermal thickness, but no change in epidermal thickness, were also noted. These results are similar to studies treating patients for other scarring conditions. An increase in collagen and elastin was seen in two studies treating patients with striae distensae (Aust et al., 2010b; Park et al., 2012), with one additionally showing an increase in epidermal thickness (Park et al., 2012). A study treating burn scars showed an increase in collagen (Aust et al., 2010a). Increases in collagen, elastin, and epidermal and dermal thickness have also been seen with skin needling in nonscarring conditions such as photodamage and skin laxity (Aust et al., 2008a, 2011; Fabbrocini et al., 2011a; Fernandes, 2005; Fernandes and Signorini, 2008; Kim et al., 2011).

Optimal number of treatments, duration between treatments, and needle length

There have been no studies directly evaluating the optimal number of treatments, time between treatments, or needle length. The median number of treatments in studies using skin needling alone was three (Alam et al., 2014; Beretta et al., 2008; Dogra et al., 2014; Fabbrocini et al., 2009, 2011c, 2014b; Leheta et al., 2011; Majid, 2009; Schwarz and Laaff, 2011; Sharad, 2011), with a range of one to five treatments. The median duration between treatments was 4 weeks, with a range of 2 to 8 weeks. The median needle length used was 1.5 mm, ranging from 1 mm to 3 mm. No studies compared needle thickness or degree of pressure applied.

Adverse events

Eighteen patients out of 246, over 10 studies (Alam et al., 2014; Beretta et al., 2008; Dogra et al., 2014; Fabbrocini et al., 2009, 2011c, 2014b; Leheta et al., 2011; Majid, 2009; Schwarz & Laaff, 2011; Sharad, 2011) treating patients with skin needling alone, had adverse events reported. Skin needling is expected to cause temporary erythema, pain, a burning sensation, edema, bleeding, or a serous ooze resolving with crusting or scabbing (Alam et al., 2014; Beretta et al., 2008; Dogra et al., 2014; Fabbrocini et al., 2009, 2011c, 2014b; Leheta et al., 2011; Majid, 2009; Sharad, 2011). Bruising and hematomas are also expected, particularly over bony prominences (Dogra et al., 2014; Fabbrocini et al., 2014b; Schwarz and Laaff, 2011). The development of “tram trek” scarring was reported with 2-mm needles in one case report (Pahwa et al., 2012). This phenomenon also occurred in a study using 1.5-mm needles (Dogra et al., 2014), but not in a study using 2.5-mm needles (Gadkari and Nayak, 2014). It is therefore unclear if this phenomenon is related to needle length. There were no other reports of scarring. Other adverse events included the development of acne and the formation of milia (Beretta et al., 2008; Leheta et al., 2011; Sharad, 2011) (see Fig. 2). Serious adverse events worth noting when skin needling was used for other dermatological conditions included facial allergic granuloma and systemic hypersensitivity reactions, possibly related to topical products put on the skin before needling or to the needles themselves (Patsou, 2013; Soltani-Arabshahi et al., 2014).

Infection as an adverse event

No studies reported bacterial infections after treatment, although some opted for topical or oral antibiotic prophylaxis (Alam et al., 2014; Gadkari and Nayak, 2014; Majid, 2009; Sharad, 2011). One study reported infections with herpes simplex virus (HSV), but it was unclear if these patients had acne scarring (Aust et al., 2008a). Reports of HSV infections have been noted in other articles (Fernandes, 2005; Torezan et al., 2013) and oral acyclovir was given to patients with a history of HSV in at least one study including patients with acne scarring (Alam et al., 2014).

Post-inflammatory hyperpigmentation as an adverse event

In an early study using rats, post-inflammatory hyperpigmentation after skin needling was found to be unlikely (Aust et al., 2008b). Of the patients treated with skin needling alone, nine developed postinflammatory hyperpigmentation (Dogra et al., 2014; Majid, 2009; Sharad, 2011); all had skin phototypes of three or greater except one, where the skin phototype was not mentioned. Not all studies reported skin types, but in at least four studies (Alam et al., 2014; Dogra et al., 2014; Fabbrocini et al., 2014b; Sharad, 2011), there were a total of 105 patients with skin phototypes of three or greater.

Conclusion

Skin needling is a relatively simple, cost-effective technique used for the treatment of acne scarring. A review of the current literature suggests that it has moderate efficacy. However, the evidence for this is limited as it is based on predominantly observational studies, which are heterogeneous in design with some lacking statistical analysis and internal validity. Skin needling was shown to work well in combination with other treatments for acne scarring, but the results of each study were specific for each treatment and lacked external validity. Skin needling appeared to be safe with a low frequency of side effects. No consensus has been reached on the use of antibiotic prophylaxis. Double-blinded randomized controlled trials are needed to further evaluate efficacy, and specific studies are needed to define the optimal number of treatments, duration between each treatment, and needle depth, and to further characterize adverse reactions.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.ijwd.2015.03.004>.

References

- Alam M, Han S, Pongpruthiphan M, Disphanurat M, Kakar R, Nodzenski M, et al. Efficacy of a needling device for the treatment of acne scars. A randomized clinical trial. *JAMA Dermatol* 2014;150(8):844–9.
- Aust MC, Fernandes D, Kolokythas P, Kaplan HM, Vogt PM. Percutaneous collagen induction therapy. An alternative treatment for scars, wrinkles, and skin laxity. *Plast Reconstr Surg* 2008;121:1421–9.
- Aust MC, Reimers K, Repenning C, Stahl F, Jahn S, Guggenheim M, et al. Percutaneous collagen induction: minimally invasive skin rejuvenation without risk of hyperpigmentation—fact or fiction? *Plast Reconstr Surg* 2008;122:1553–63.
- Aust MC, Knobloch K, Reimers K, Redeker J, Ipaktchi R, Altintas MA, et al. Percutaneous collagen induction therapy: An alternative treatment for burn scars. *Burns* 2010;36:836–43.
- Aust MC, Knobloch K, Vogt PM. Percutaneous collagen induction as a novel therapeutic option for Striae Distensae. *Plast Reconstr Surg* 2010;126:4.
- Aust MC, Reimers K, Kaplan HM, Stahl F, Repenning C, Scheper T, et al. Percutaneous collagen induction regeneration in place of cicatrization? *J Plast Reconstr Aesthet Surg* 2011;64:97e–107e.
- Badran MM, Kuntsche J, Fahr A. Skin penetration enhancement by a microneedle device (Dermaroller) in vitro: dependency on needle size and applied formulation. *Eur J Pharm Sci* 2009;36:511–23.
- Baryza MJ. The Vancouver Scar Scale: an administration tool and its interrater reliability. *J Burn Care Rehabil* 1995;16:535.
- Bencini PL, Galimberti MG, Pellacani G, Longo C. Application of photodynamic therapy combined with pre-illumination microneedling in the treatment of actinic keratosis in organ transplant recipients. *Br J Dermatol* 2012;167:1193–4.
- Beretta D, Clementoni MT, Pinelli S, Signorini M. Induzione percutanea del collagene (needling): valutazione critica. *J Plast Dermatol* 2008;4(3):291–300.
- Budamakuntla L, Loganathan E, Suresh DH, Shanmugam S, Suryanarayan S, Dongare A, et al. Open-label, comparative study of tranexamic acid microinjections and tranexamic acid with microneedling in patients with melasma. *J Cutan Aesthet Surg* 2013;6(3):139–43.
- Camirand A, Doucet J. Needle dermabrasion. *Aesthet Plast Surg* 1997;21:48–51.
- Clementoni MT, B-Roscher M, Munavalli GS. Photodynamic photorejuvenation of the face with a combination of microneedling, red light, and broadband pulsed light. *Lasers Surg Med* 2010;42:150–9.
- Doddaballapur S. Microneedling with dermaroller. *J Cutan Aesthet Surg* 2009;2(2):110–1.
- Dogra S, Yadav S, Sarangal R. Microneedling for acne scars in Asian skin type: an effective low cost treatment modality. *J Cosmet Dermatol* 2014;13:180–7.
- Dreno B, Khammari A, Orain N, Noray C, Mériat-Kieny C, Méry S, et al. ECCA grading scale: an original validated acne scar grading scale for clinical practice in dermatology. *Dermatology* 2007;214(1):46–51.
- Fabbrocini G, Fardella N, Monfrecola A, Proietti I, Innocenzi D. Acne scarring treatment using skin needling. *Clin Exp Dermatol* 2009;34:874–9.
- Fabbrocini G, De Vita V, Di Costanzo L, Francesco P, Mauriello MC, Monfrecola A, et al. Skin needling in the treatment of the aging neck. *Skinmed* 2011;9:347–51.
- Fabbrocini G, De Vita V, Fardella N, Pastore F, Annunziata MC, Mauriello MC, et al. Skin needling to enhance depigmenting serum penetration in the treatment of melasma. *Plast Surg Int* 2011;2011:158241.
- Fabbrocini G, De Vita V, Pastore F, Panariello L, Fardella N, Sepulveres R, et al. Combined use of skin needling and platelet-rich plasma in acne scarring treatment. *Cosmet Dermatol* 2011;24(4):177–83.
- Fabbrocini G, De Vita V, Izzo R, Monfrecola G. The use of skin needling for the delivery of a eutectic mixture of local anesthetics. *G Ital Dermatol Venereol* 2014;149(5):581–5.
- Fabbrocini G, De Vita V, Monfrecola A, Pia De Padova M, Brazzini B, Teixeira F, et al. Percutaneous collagen induction: an effective and safe treatment for post-acne scarring in different skin phototypes. *J Dermatolog Treat* 2014;25:147–52.
- Fernandes D. Minimally invasive percutaneous collagen induction. *Oral Maxillofac Surg Clin North Am* 2005;17(1):51–63 (vi).

- Fernandes D, Signorini M. Combating photoaging with percutaneous collagen induction. *Clin Dermatol* 2008;26:192–9.
- Gadkari R, Nayak C. A split-face comparative study to evaluate efficacy of combined subcision and dermaroller against combined subcision and cryoroller in treatment of acne scars. *J Cosmet Dermatol* 2014;13:8–43.
- Garg S, Baveja S. Combination therapy in the management of atrophic acne scars. *Cutan Aesthet Surg* 2014;7(1):18–23.
- Goodman GJ, Baron JA. Postacne scarring: a qualitative global scarring grading system. *Dermatol Surg* 2006;32:1458–66.
- Goodman GJ, Baron JA. Postacne scarring: a quantitative global scarring grading system. *J Cosmet Dermatol* 2006;5:48–52.
- Kalluri H, Kolli CS, Banga AK. Characterization of microchannels created by metal microneedles: formation and closure. *AAPS J* 2011;13(3):473–81.
- Kang H, Son T, Yoon J, Kwon K, Nelson JS, Jung B. Evaluation of laser beam profile in soft tissue due to compression, glycerol, and micro-needling. *Lasers Surg Med* 2008;40:570–5.
- Kang WH, Kim YJ, Pyo WS, Park SJ, Kim JH. Atrophic acne scar treatment using triple combination therapy: dot peeling, subcision and fractional laser. *J Cosmet Laser Ther* 2009;11:212–5.
- Kim S. Pinpoint irradiation with CO₂ laser and needling with a 26 gauge needle. Elicina cream for a few days after each treatment. *J Cosmet Laser Ther* 2008;10:177–80.
- Kim S, Lee J, Kwon HB, Ahn B, Lee A. Greater collagen deposition with the microneedle therapy system than with intense pulsed light. *Dermatol Surg* 2011;37:336–41.
- Lee JB, Chung WG, Kwahck H, Lee KH. Focal treatment of acne scars with trichloroacetic acid: chemical reconstruction of skin scars method. *Dermatol Surg* 2002;28:1017–21.
- Leheta TM, Tawdy AE, Hay RA, Farid S. Percutaneous collagen induction versus full-concentration trichloroacetic acid in the treatment of atrophic acne scars. *Dermatol Surg* 2011;37:207–16.
- Leheta TM, Abdel Hay RM, El Garem YF. Deep peeling using phenol versus percutaneous collagen induction combined with trichloroacetic acid 20% in atrophic post-acne scars; a randomized controlled trial. *J Dermatolog Treat* 2014;25:130–6.
- Leheta TM, Abdel Hay RM, Hegazy RA, El Garem YF. Do combined alternating sessions of 1540 nm nonablative fractional laser and percutaneous collagen induction with trichloroacetic acid 20% show better results than each individual modality in the treatment of atrophic acne scars? A randomized controlled trial. *J Dermatolog Treat* 2014;25:137–41.
- Lipper GM, Perez M. Nonablative acne scar reduction after a series of treatments with a short-pulsed 1,064-nm neodymium: YAG laser. *Dermatol Surg* 2006;32:998–1006.
- Majid I. Microneedling therapy in atrophic facial scars: an objective assessment. *J Cutan Aesthet Surg* 2009;2:26–30.
- Mohammed G. Randomized clinical trial of CO₂ laser pinpoint irradiation technique with/without needling for ice pick acne scar. *J Cosmet Laser Ther* 2013;15:177–82.
- Nofal E, Helmy A, Nofal A, Alakad R, Nasr M. Platelet-rich plasma versus CROSS technique with 100% trichloroacetic acid versus combined skin needling and platelet-rich plasma in the treatment of atrophic acne scars: a comparative study. *Dermatol Surg* 2014;40:864–73.
- Orentreich DS, Orentreich N. Subcutaneous incisionless (subcision) surgery for the correction of depressed scars and wrinkles. *Dermatol Surg* 1995;21:543–9.
- Pahwa M, Pahwa P, Zaheer A. “Tram track effect” after treatment of acne scars using a microneedling device. *Dermatol Surg* 2012;38:1107–8.
- Park KY, Kim HK, Kim SE, Kim BJ, Kim MN. Treatment of striae distensae using needling therapy: a pilot study. *Dermatol Surg* 2012;38:1823–8.
- Patsou P. Severe systemic reaction associated with skin microneedling therapy in two sisters: a previously unrecognized potential for complications? *JAAD* 2013;68(4):AB219.
- Peterson JD, Palm MD, Kiripolsky MG, Guiha IC, Goldman MP. Evaluation of the effect of fractional laser with radiofrequency and fractionated radiofrequency on the improvement of acne scars. *Dermatol Surg* 2011;37:1260–7.
- Schwarz M, Laaff H. A prospective controlled assessment of microneedling with the dermaroller device. *Plast Reconstr Surg* 2011;127(6):146e–8e.
- Sharad J. Combination of microneedling and glycolic acid peels for the treatment of acne scars in dark skin. *J Cosmet Dermatol* 2011;10:317–23.
- Soltani-Arabshahi R, Wong JW, Duffy KL, Powell DL. Facial allergic granulomatous reaction and systemic hypersensitivity associated with microneedle therapy for skin rejuvenation. *JAMA Dermatol* 2014;150(1):68–72.
- Torezan L, Chaves Y, Niwa A, Sanches JA, Festa-Neto C, Szeimies R. A pilot split-face study comparing conventional methyl aminolevulinic acid-photodynamic therapy (PDT) with microneedling-assisted PDT on actinically damaged skin. *Dermatol Surg* 2013;39:1197–201.
- Yoon J, Son T, Choi E, Choi B, Nelson JS, Jung B. Enhancement of optical skin clearing efficacy using a microneedle roller. *J Biomed Opt* 2008;13(2):021103.
- Yoon J, Park D, Son T, Seo J, Nelson JS, Jung B. A physical method to enhance transdermal delivery of a tissue optical clearing agent: combination of microneedling and sonophoresis. *Lasers Surg Med* 2010;42:412–7.